

Pion gas viscosity at low temperature and density.

Antonio Dobado*

*Departamento de Física Teórica,
Universidad Complutense de Madrid, 28040 Madrid, Spain*

Silvia N. Santalla†

*Departamento de Física,
Universidad Carlos III de Madrid, 28911 Leganés, Spain*

By using Chiral Perturbation Theory and the Uehling-Uhlenbeck equation we compute the viscosity of a pion gas, in the low temperature and low density regime, in terms of the temperature, and the pion fugacity. The viscosity turns out to be proportional to the squared root of the temperature over the pion mass. Next to leading corrections are proportional to the temperature over the pion mass to the 3/2.

PACS numbers: 11.30.Rd, 12.39.Fe, 24.10.Nz, 25.75.-q

I. INTRODUCTION

The possibility of discovering the so called quark-gluon plasma at RHIC (Relativistic Heavy Ion Collider) or in the future LHC (Large Hadron Collider) has triggered a lot of effort in the theoretical description of ultra-relativistic heavy ion collisions. The modern view of these collisions was set mainly by Bjorken [1] and it is based in the hydrodynamic model already considered by Landau [2]. In the last stages of the evolution of the central rapidity region the hadronic fluid is made mainly of pions. It has been argued that this expanding pion gas can reach thermal equilibrium much more efficiently than chemical equilibrium since at low energies pion interactions are mostly elastic [3]. In order to have an appropriate phenomenological description of the expanding pion gas it is needed to set the proper hydrodynamic equations. In the pioneering work by Bjorken the hadronic gas (both in the confined and in the quark-gluon plasma phases) was assumed to be a perfect relativistic fluid and accordingly viscosity and heat conductivity were neglected. In order to check the validity of this assumption in the different regimes of the pion gas expansion it is needed to compute the corresponding transport coefficients as a function of the temperature and density. This could make possible to know in what physical situations they are relevant and the ideal fluid equations are not appropriate any more. In addition the expressions obtained for these coefficients can be used to set the correct relativistic version of the Navier-Stokes equations which applies when dissipative effects must be taken into account in the pion fluid dynamics. Curiously enough, the computation of these coefficients is also interesting from the fundamental point of view, since it can be done completely from first principles.

In this work we will illustrate this by computing the viscosity of the pion gas at low energy and density. Our computation will rely just in standard kinetic theory and the chiral symmetry properties of the strong interaction and thus our results can be considered model independent. In fact our results will depend only on the pion mass M_π and the pion decay constant F_π . The main points on which our computation is based are the following. First of all our computation is going to apply only at low temperature and density. At low temperatures hadronic matter is confined into hadrons and at very low temperatures the only excited modes are pions for vanishing barionic number n_B density. Notice that $n_B \sim 0$ is supposed to be the case in the central rapidity region of heavy ion collisions. Moreover the additional low-density condition allows us to consider the non barionic hadronic matter as a pion gas.

At low temperatures most of the pion interactions occur at low energies. The low energy pion interactions are completely determined by the chiral symmetry pattern of the strong interactions. In the chiral limit (vanishing u and d quark masses) the strong interaction dynamics is invariant under the $SU(2)_L \times SU(2)_R$ chiral transformations. However this symmetry is supposed to be spontaneously broken to the diagonal or isospin group $SU(2)_{L+R}$, being the pions the corresponding Goldstone bosons. In the real world quarks masses are different from zero and then this symmetry breaking pattern is not exact resulting in a small value (when compared with other hadrons) for the pion mass M_π . In any case chiral symmetry determines completely the low energy pion interactions in terms of M_π and F_π (Weinberg low energy theorems). The corrections appearing at higher energies can be computed in a systematic way by means of Chiral Perturbation Theory [4]. In the chiral limit the pion scattering amplitude goes to zero in the low energy limit. Out of the chiral limit it is proportional to M_π^2/F_π^2 . In any case the interaction is small in this regime. Therefore we arrive to the conclusion that non barionic hadronic matter at low temperature and density can be described as

*Electronic address: malcon@fis.ucm.es

†Electronic address: ssantall@fis.uc3m.es

a weakly interacting pion gas. Another important observation is that at low temperatures (energies) most of the pion collisions are elastic. This fact implies in particular that the number of pions is conserved. Thus it is possible to introduce the corresponding pion chemical potential μ (not to be confused with the more usual chemical potential associated to the barionic number). This makes it possible to consider the pion gas at thermal equilibrium at temperature T for different values of the chemical potential μ , or what it is the same, different pion densities, and not only the case of chemical equilibrium corresponding to $\mu = 0$.

According to the previous discussion the simplest state for non barionic hadronic matter at low temperatures and densities is just a gas of weakly interacting pions in thermal equilibrium which could be described in a first approximation with a Bose-Einstein distribution. In addition, at very low temperatures the average pion energy is much smaller than the pion mass allowing for to have a non relativistic description of the pion gas. As discussed later, the low density condition will prevent the formation of the Bose-Einstein condensate out of the pions, thus making the description of the hadronic matter in the mentioned regime especially simple.

More generally it is possible to consider the case when the equilibrium is only local. In this case the temperature, density (or pion chemical potential) and the overall velocity are space-time functions. In this situation it is possible to have an hydrodynamic description of the system as that considered in the Bjorken picture of heavy ion collisions. A departure from local equilibrium will give rise to dissipative processes like viscosity, which is the main topic of this work. The evolution of the distribution functions outside equilibrium can be studied by using the Uehling-Uhlenbeck equation, which is the quantum version of the Boltzmann transport equation. The computation of the transport coefficients such as viscosity or heat conductivity requires the solution of the transport equation for different specific perturbations of the local equilibrium distribution.

As it was mentioned above, in this work we are interested in the computation of the viscosity of the hadronic fluid with the approximations discussed in the previous paragraphs. In fact it is possible to have a good estimation of this magnitude by neglecting quantum effects. In this case the equilibrium distribution function is just the Boltzmann distribution instead the Bose-Einstein distribution. Moreover the elastic scattering pion cross section is constant at low energies and as we will see it is given by

$$\sigma = \frac{23}{384} \frac{M_\pi^2}{\pi F_\pi^4} \quad (1)$$

In addition, at low temperatures we can use the non relativistic approximation so that our problem is equivalent to the classical computation of the viscosity of a gas of hard spheres of radius R so that $\sigma = \pi R^2$. By using the well known result in this case (see for instance [5]) we

find the viscosity to be given by

$$\eta = \frac{1920}{368} \frac{F_\pi^4 \pi^{3/2}}{M_\pi} \sqrt{\frac{T}{M_\pi}} \quad (2)$$

This formula provides a nice estimation of the viscosity of the pion gas. However it does not take into account any quantum effect that can be important at low temperatures. In addition it does not include any dependence in the pion chemical potential, i.e. in the pion density.

In the following we will show how these effects can be properly taken into account to find the pion gas viscosity in terms of the temperature and chemical potential (or density) in the low temperature and density regime at the leading order. We will also estimate the magnitude of the next to leading corrections. The plan of the paper goes as follows: In sec.2 we review the main equations of the non-relativistic ideal pion gas to set the notation and for further reference. In sec.3 we describe briefly the fundamentals of the kinetic theory needed to follow the present work and in sec.4 we show how the hydrodynamic equations emerge from it. In sec.5 we obtain the equations for the transport coefficients in terms of the appropriate solutions of the linearized transport equations. In sec.6 we simplify these equations to write them in useful form for finding explicit formulae for the viscosity. In sec.7 we give the cross-section for the low energy elastic pion scattering obtained from the lowest order chiral perturbation theory. In sec.8 we obtain the lowest order terms of the pion gas viscosity. In sec.9 we show our numerical results and discuss the magnitude of the next to leading corrections and in sec.10 we set the main conclusions of our work. Finally in the appendix we study the properties of the polynomials introduced in this work that play a role similar to the standard Sonine polynomials in the more familiar classic computations.

II. STATE EQUATION

In this section we review briefly the equation of state of a free non-relativistic bosonic gas, which will provide our description of the pion gas at low temperature and density. As it is well known [7], for a large volume the pressure can be written as

$$P = - \frac{gT}{2\pi^2} \int_0^\infty dp p^2 \log [1 - e^{\beta(\mu - E(p))}] - \frac{gT}{V} \log [1 - e^{\beta\mu}] \quad (3)$$

where g is the number of pions ($g = 3$), $T = 1/\beta$ is the temperature, V is the volume, μ is the pion chemical potential and $E(p)$ is the energy of a non relativistic pions in terms of the momentum i.e. $\vec{p} = M_\pi \vec{v}$, $E(p) = p^2/2M_\pi$. The number density or the number of pions per unit of volume is given by

$$n = \frac{g}{2\pi^2} \int_0^\infty dp p^2 \frac{1}{e^{-\beta(\mu - E(p))} - 1} + \frac{g}{V} \frac{1}{e^{-\beta\mu} - 1} \quad (4)$$

where the last term is the number of pions with zero momentum per unit of volume. When this term is not negligible it is responsible for the Bose-Einstein condensate. We thus define

$$n_0 = \frac{g}{V} \frac{1}{e^{-\beta\mu} - 1} \quad (5)$$

Clearly the above equations make sense only when the chemical potential satisfies $\mu \leq 0$. As a consequence, in the thermodynamic limit, where $N, V \rightarrow \infty$ with N/V constant, we find two phases: In the first one (normal phase) $\mu \leq 0$ and $n_0 = 0$. However if we lower the temperature, keeping a fixed density, μ increases until $\mu = 0$. At that point $n_0 \neq 0$ and the ground state density starts to grow forming the Bose-Einstein condensate. Eventually, at $T = 0$, all pions are in the ground state so that $n = n_0$. The critical temperature T_c where the phase transition occurs can be found to be

$$T_c \equiv \frac{2\pi}{M_\pi} \left(\frac{n}{g\zeta_{3/2}} \right)^{2/3}$$

where $\zeta_{3/2}$ is the Riemann ζ function evaluated at $3/2$. Therefore the critical temperature can be made arbitrary small by lowering the density. In this work we are interested in the the low temperature and the low density regime. In order to simplify the analysis we will not consider the contribution of the condensate. Thus for a given temperature T , we will always assume that the density will be low enough so that $T > T_c$ and no significant fraction of pions will have zero momentum (see [6] for a discussion of the pion condensate).

In the following it will also be convenient to introduce the fugacity

$$z \equiv \exp\left(\frac{\mu}{T}\right)$$

and then the above equations for the density and the pressure can be written as

$$n = \frac{g}{2\pi^2} \int_0^\infty \frac{p^2}{z^{-1}e^{\frac{p^2}{2M_\pi T}} - 1} dp \quad (6)$$

and

$$P = -\frac{gT}{2\pi^2} \int_0^\infty p^2 \ln \left(1 - ze^{-\frac{p^2}{2M_\pi T}} \right) dp \quad (7)$$

Introducing the variable $x = p^2/(2M_\pi T)$ we have

$$n = \frac{g}{4\pi^2} (2M_\pi T)^{3/2} \int_0^\infty \frac{x^{1/2}}{z^{-1}e^x - 1} dx \quad (8)$$

and

$$P = \frac{g}{6\pi^2} (2M_\pi T)^{3/2} T \int_0^\infty \frac{x^{3/2}}{z^{-1}e^x - 1} dx \quad (9)$$

where the integrals can be written in terms of the g_n functions [8] as

$$\int \frac{x^n}{z^{-1}e^x - 1} dx = \Gamma(n+1)g_{n+1}(z) \quad (10)$$

with

$$g_n(z) = \sum_{l=1}^{\infty} \frac{z^l}{l^{n+1}}$$

so that

$$g_s(1) = \zeta_s$$

and

$$g_{s-1}(z) = z \frac{dg_s(z)}{dz}$$

Therefore we have for a non relativistic free pion gas at equilibrium at low temperature and density

$$n = g \left(\frac{M_\pi T}{2\pi} \right)^{3/2} g_{3/2}(z) \quad (11)$$

and

$$P = g \left(\frac{M_\pi T}{2\pi} \right)^{3/2} T g_{5/2}(z) \quad (12)$$

Thus the equation of state can be written as

$$P = nT \frac{g_{5/2}(z)}{g_{3/2}(z)} \quad (13)$$

III. THE KINETIC THEORY FOR PIONS

The possibility of having an hydrodynamic description of the pion gas requires the definition of macroscopic fields such as the velocity $\vec{V} = \vec{V}(\vec{r}, t)$, the pressure $P = P(\vec{r}, t)$, the number density $n = n(\vec{r}, t)$, the temperature $T = T(\vec{r}, t)$ and others. In other words, it is needed to have a large number of pions in each volume element dV . The statistical description of the gas is based on the (one-particle) distribution function $f(\vec{r}, \vec{v}, t)$ which gives the number of pions inside the phase-space volume element $d\vec{r} d\vec{v}$ at the instant t . In fact the phase-space volume element is given by $d\vec{r} d\vec{p}$ but in the case considered here the difference amounts to a constant factor which we include in the definition of the distribution function f . Thus the total number of pions will be given by

$$N = \int f(\vec{r}, \vec{v}, t) d\vec{r} d\vec{v} = \int n(\vec{r}, t) d\vec{r} \quad (14)$$

As we are interested here in the behaviour of the pion gas at low temperatures and densities, most of the the pion interactions will occur at low energies. Then the pion-pion interactions will be essentially elastic and the

total number of pions will be a conserved quantity. This in particular means that it makes sense to introduce the chemical potential $\mu = \mu(\vec{r}, t)$ related with the pion number even in presence of low-energy pion interactions. The evolution of the distribution function is determined by the BBKGY hierarchy (from N.N. Bogoliubov, M. Born, G. Kirkwood, H.S. Green and J. Yvon) [9]. This is a set of N coupled equations for the n -particle distribution functions $f_{(n)}(\vec{r}_1, \vec{v}_1, \vec{r}_2, \vec{v}_2, \dots, \vec{r}_n, \vec{v}_n, t)$ with $f = f_{(1)}$. However, for low-density gases it can be a good approximation to truncate the BBKGY hierarchy. For example by using the Bogoliubov ansatz, i.e. by writing the two-particle function in terms of an appropriate product of two one-particle functions (molecular chaos hypothesis), it is possible to decouple the first BBKGY equation from the others to obtain the well known Boltzmann equation. This equation was modified by Uehling and Uhlenbeck in order to include the quantum statistic effects (see for example [10]). For the bosonic case the Uehling-Uhlenbeck equation reads

$$(\partial_t + \vec{v}_1 \vec{\nabla}_{\vec{r}}) f_1 = C[f_1] \quad (15)$$

where

$$C[f_1] = \int d\vec{v}_2 d\sigma u \left[f'_1 f'_2 \left(1 + \frac{f_1}{A}\right) \left(1 + \frac{f_2}{A}\right) - f_1 f_2 \left(1 + \frac{f'_1}{A}\right) \left(1 + \frac{f'_2}{A}\right) \right] \quad (16)$$

is the collision term or collision functional. The above equation describes the irreversible space-time evolution of the (one-point) distribution function f , being this irreversible behaviour a consequence of the truncation of the BBKGY hierarchy. As usual we use for short the notation $f_i = f(\vec{r}_i, \vec{v}_i, t)$ and $f'_i = f(\vec{r}'_i, \vec{v}'_i, t)$ with $i = 1, 2$. $d\sigma$ is the differential cross-section corresponding to the elastic scattering of two pions with initial and final velocities $\vec{v}_1, \vec{v}_2$ and $\vec{v}'_1, \vec{v}'_2$, respectively. u is the relative velocity of the initial particles $u = |\vec{v}_1 - \vec{v}_2|$ and, finally, the normalization constant A is given by $A = gM_\pi^3 / (2\pi)^3$.

As it is the case of the Boltzmann equation, the Uehling-Uhlenbeck equation typically drives the gas from arbitrary initial distributions to the equilibrium distribution f^0 which is a fixed point of this equation in the sense

that the collision functional vanishes on it

$$C[f^0] = 0 \quad (17)$$

For a gas moving at a macroscopic velocity $\vec{V}$ the equilibrium distribution corresponding to temperature T and pion chemical potential μ is given by

$$f^0(\vec{r}, \vec{v}, t) = A \left(e^{\frac{1}{T} \left(M_\pi \frac{(\vec{v} - \vec{V})^2}{2} - \mu \right)} - 1 \right)^{-1} \quad (18)$$

from which the properties of free pion gas given in the previous section can be derived. The case of local equilibrium is also described by this distribution but having the $\vec{V}, T$ and μ parameters depending on the position and time. This case corresponds to an ideal fluid. However, in order to describe dissipative processes, such as viscosity or thermal conduction, it is needed to consider the case where the pion gas is not even in local thermal equilibrium.

IV. HYDRODYNAMIC EQUATIONS

Given some magnitude ψ it is possible to compute its statistic average by using the distribution function as

$$\langle \psi \rangle = \frac{\int f \psi d\vec{r}}{\int f d\vec{r}} = \frac{\int f \psi d\vec{r}}{n(\vec{r}, t)} \quad (19)$$

where $n = n(\vec{r}, t)$ is the pion number density. This average can be understood as the corresponding macroscopic magnitude in the hydrodynamic approach [11]. For example, the macroscopic fluid velocity $\vec{V}$ is given by

$$\vec{V} = \frac{\int f \vec{v} d\vec{v}}{\int f d\vec{v}} = \frac{1}{n} \int f \vec{v} d\vec{v} \quad (20)$$

The equation governing the evolution of any of these magnitudes (transport equations) can be obtained by multiplying ψ_1 by the Uehling-Uhlenbeck equation and then integrating $\vec{v}_1$ so that

$$\int d\vec{v}_1 (\partial_t + \vec{v}_1 \vec{\nabla}_{\vec{r}}) f_1 \psi_1 = \int d\vec{v}_1 d\vec{v}_2 d\sigma u \psi_1 (f'_1 f'_2 (1 + A^{-1} f_1) (1 + A^{-1} f_2) - f_1 f_2 (1 + A^{-1} f'_1) (1 + A^{-1} f'_2)) \quad (21)$$

The r.h.s. term can be written in a more symmetric fashion as

$$\frac{1}{4} \int d\vec{v}_1 d\vec{v}_2 d\vec{v}_3 d\vec{v}_4 |T|^2 (\psi_1 + \psi_2 - \psi'_1 - \psi'_2) (f'_1 f'_2 (1 + A^{-1} f_1) (1 + A^{-1} f_2) - f_1 f_2 (1 + A^{-1} f'_1) (1 + A^{-1} f'_2)) \quad (22)$$

where $|T|^2$ is defined through the relation

$$d\sigma = \frac{|T|^2}{|\vec{v}_1 - \vec{v}_2|} d\vec{v}'_1 d\vec{v}'_2 \quad (23)$$

From this form of the collision terms it becomes apparent than it vanishes for any quantity ψ which is conserved

in the collision such as the total energy or momentum. On the other hand, by using the definition of statistical average, the first term of (21) can be written as:

$$\partial_t n \langle \psi \rangle + \vec{\nabla} n \langle \psi \vec{v} \rangle \quad (24)$$

Thus for conserved magnitudes this term vanishes. In order to find the basic hydrodynamic equations it is useful

to write the velocity as $\vec{v} = \vec{V} + \vec{\xi}$, i.e. $\vec{\xi}$ is the pion velocity relative to the macroscopic velocity $\vec{V}$ and therefore it averages to zero. Choosing $\psi = 1$, $\vec{v}$ and v^2 (corresponding to the conservation of a constant, the momentum and the energy in the elastic collision) we obtain the equations

$$\begin{aligned} \psi = 1 &\longrightarrow \partial_t n + \vec{\nabla} n \vec{V} = 0 \\ \psi = \vec{v} &\longrightarrow \partial_t n V_i + \nabla_j n V_i V_j + \nabla_j n \langle \xi_i \xi_j \rangle = 0 \\ \psi = v^2 &\longrightarrow \partial_t n (V^2 + \langle \xi^2 \rangle) + \nabla_i n V_i (V^2 + \langle \xi^2 \rangle) + \nabla_i 2n V_j \langle \xi_i \xi_j \rangle + \nabla_i n \langle \xi_i \xi^2 \rangle = 0 \end{aligned} \quad (25)$$

In order to understand the meaning of the above equations we can introduce the following macroscopic quantities:

$$P_{ij} = M_\pi n \langle \xi_i \xi_j \rangle \quad (26)$$

is the pressure tensor,

$$Q = \frac{1}{2} M_\pi n \langle \xi^2 \rangle \quad (27)$$

is the internal energy density and

$$q_i = \frac{1}{2} M_\pi n \langle \xi_i \xi^2 \rangle \quad (28)$$

is the energy (heat) flux. Due to the isotropy of the pion fluid the pressure tensor can be taken as diagonal, i.e. $P_{ij} = P \delta_{ij}$. In this case $Q = 3P/2$ and $q_i = 0$. Thus, (by taking $\vec{V} = 0$ but not its derivatives, which can always be done) we arrive to the standard ideal fluid equations, namely the continuity equation

$$\partial_t n + n \nabla_i V_i = 0, \quad (29)$$

the Euler equation

$$n M_\pi \partial_t V_i + \nabla_i P = 0 \quad (30)$$

and the energy conservation equation

$$\partial_t Q + \frac{5}{3} Q \nabla_i V_i = 0 \quad (31)$$

In spite of the above equations for the ideal fluid, it is well established that an irreversible flux of energy and momentum appear in an inhomogeneous gas. This flux gives rise to the well known dissipative processes of thermal and conduction and viscosity. For example an small gradient of temperature produces a heat flow which in a first approximation can be written as

$$q_i = -\kappa \nabla_i T \quad (32)$$

where κ is a coefficient called thermal conductivity. In order to include momentum diffusion effects we add an extra term to the isotropic pressure tensor:

$$P_{ij} = P \delta_{ij} + P'_{ij} \quad (33)$$

The requirement for this tensor to vanish for uniform translations and rotations of the fluid, makes it possible to write it to first order in the velocity derivatives as

$$P'_{ij} = -\eta \left(\frac{\partial V_i}{\partial x_j} + \frac{\partial V_j}{\partial x_i} - \frac{2}{3} \sum_{l=1}^3 \frac{\partial V_l}{\partial x_l} \delta_{ij} \right) - \zeta \sum_{l=1}^3 \frac{\partial V_l}{\partial x_l} \delta_{ij} \quad (34)$$

Thus it is needed to introduce two coefficients η and ζ which are usually called first and second viscosity coefficients. The introduction of the dissipative effects does not modify the continuity equation. However the Euler equation becomes

$$\partial_t n V_i + \nabla_j n V_i V_j + \frac{1}{M_\pi} \nabla_j P \delta_{ij} = \frac{1}{M_\pi} \nabla_j \left[\eta \left(\frac{\partial V_i}{\partial x_j} + \frac{\partial V_j}{\partial x_i} - \frac{2}{3} \sum_{l=1}^3 \frac{\partial V_l}{\partial x_l} \delta_{ij} \right) + \zeta \sum_{l=1}^3 \frac{\partial V_l}{\partial x_l} \delta_{ij} \right] \quad (35)$$

which is known as the Navier-Stokes equation. The energy conservation equation is in this case

$$\begin{aligned} \partial_t \left(nV^2 + \frac{2}{M_\pi} Q \right) + \nabla_i V_i \left(nV^2 + \frac{2}{M_\pi} Q \right) + \nabla_i \frac{2}{M_\pi} V_j P \delta_{ij} = \\ \nabla_i \kappa \nabla_i T + \frac{2}{M_\pi} \nabla_i V_j \left[\eta \left(\frac{\partial V_i}{\partial x_j} + \frac{\partial V_j}{\partial x_i} - \frac{2}{3} \sum_{l=1}^3 \frac{\partial V_l}{\partial x_l} \delta_{ij} \right) + \zeta \sum_{l=1}^3 \frac{\partial V_l}{\partial x_l} \delta_{ij} \right] \end{aligned} \quad (36)$$

Thus the description of the dissipative flow of energy and momentum requires the introduction of three transport coefficients κ, η and ζ . However, for an incompressible fluid the second (shear) viscosity does not play any role. In the following we will assume that this is the case for the pion gas considered here.

V. COMPUTATION OF THE TRANSPORT COEFFICIENTS

As it has been mentioned above the description of the dissipative processes requires going beyond local equilibrium. Thus, in order to compute the transport coefficients we have to consider a distribution function f slightly perturbed from the equilibrium distribution f^0 , i.e.

$$f = f^0 + \delta f = f^0 + f^0 \frac{\chi}{T} \quad (37)$$

where χ is an arbitrary function of the velocities which represents the inhomogeneous contribution to the distribution function. This contribution will be assumed to be small in the sense that $\chi/T \ll 1$, since we are interested only in the computation of the transport coefficients. The χ function must be determined by solving the Uehling-Uhlenbeck equation which can be linearized with respect to χ . Then it is possible to compute the transport coefficients from this function. For example, the heat flux can be written as

$$q_i = \frac{1}{2} M_\pi \int f^0 \left(1 + \frac{\chi}{T} \right) v_i v^2 d\vec{v} \quad (38)$$

Taking into account that this flux vanishes for an homogeneous gas and using the definition of the heat conductivity we have

$$-\kappa \nabla_i T = \frac{M_\pi}{2T} \int f^0 \chi v_i v^2 d\vec{v} \quad (39)$$

Now it is useful to write the χ function as

$$\chi = -\nabla_i T h_i(\vec{v}) = -\nabla_i T v_i h(v) \quad (40)$$

Therefore

$$\kappa \nabla_i T = \kappa_{ij} \nabla_j T \quad (41)$$

with the tensor conductivity κ_{ij} being given by

$$\kappa_{ij} \equiv \frac{M_\pi}{2T} \int f^0 h v_j v_i v^2 d\vec{v} \quad (42)$$

For an isotropic gas $\kappa_{ij} = \kappa \delta_{ij}$ so that, we have the following equation for the heat conductivity

$$\kappa = \frac{M_\pi}{6T} \int f^0 h v^4 d\vec{v} \quad (43)$$

In a similar way it is possible to write the viscosity in terms of an h function. In order to do that we write the pressure tensor in terms of the distribution function

$$P_{ij} = M_\pi \int f^0 \left(1 + \frac{\chi}{T} \right) v_i v_j d\vec{v} \quad (44)$$

For the present calculation it is enough to consider the case $\vec{V} = 0$ (but not its derivatives). Then the non isotropic component of the pressure tensor can be written as

$$P'_{ij} = \frac{M_\pi}{T} \int f^0 \chi v_i v_j d\vec{v} \quad (45)$$

On the other hand for an incompressible fluid this tensor is

$$P'_{ij} = -2\eta \left(V_{ij} - \frac{1}{3} \nabla_h V_h \delta_{ij} \right) \quad (46)$$

where we have introduced $V_{ij} = (\partial_i V_j + \partial_j V_i)/2$. For further convenience we choose the perturbation to have the form

$$\chi = - \left(V_{kl} - \frac{1}{3} \nabla_h V_h \delta_{kl} \right) h_{kl}(\vec{v}) \quad (47)$$

where h_{kl} is a microscopic velocity dependent quantity which can be written as

$$h_{kl} = \left(v_k v_l - \frac{1}{3} v^2 \delta_{kl} \right) h(v) \quad (48)$$

Therefore, by equating the two forms of P'_{ij} we find

$$2\eta \left(V_{ij} - \frac{1}{3} \nabla_h V_h \delta_{ij} \right) = \eta_{ijkl} \left(V_{kl} - \frac{1}{3} \nabla_h V_h \delta_{kl} \right) \quad (49)$$

where η_{ijkl} (the viscosity tensor) is given by

$$\eta_{ijkl} \equiv M_\pi T \int v_i v_j \left(v_k v_l - \frac{1}{3} v^2 \delta_{kl} \right) f^0(v) h(v) d\vec{v} \quad (50)$$

Due to the several symmetries of this tensor it can be written as

$$\eta_{ijkl} = \eta \left(\delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk} - \frac{2}{3}\delta_{ij}\delta_{kl} \right) \quad (51)$$

so that we have

$$\eta = \frac{M_\pi}{10T} \int v_i v_j \left(v_i v_j - \frac{1}{3} v^2 \delta_{ij} \right) f^0(v) h(v) d\vec{v} \quad (52)$$

As the h function depends only on the velocity modulus it is possible to perform the angular integrations

$$\eta = \frac{4\pi}{15T} A \int_0^\infty \frac{1}{z^{-1} \exp\left(\frac{M_\pi v^2}{2T} - 1\right)} v^6 h(v) d\vec{v} \quad (53)$$

In the following it will be useful to consider h as a function of the adimensional variable x , i.e. $h = h(x)$ defined as

$$x = \frac{M_\pi v^2}{2T} \quad (54)$$

At this point it is customary to develop the $h(x)$ in terms of the Sonine polynomials. However it is more appropriate in our case to introduce a new family of orthogonal polynomials $P_r^s(z; x)$ since we are considering the Uehling-Uhlenbeck as the transport equation (i.e. we have taken into account quantum effects) instead of the more common Boltzmann equation. The definition and the main properties of this new polynomials can be found in appendix A. In terms of P_r^s the h function can be written as

$$h(x) = \sum_{s=0}^{\infty} B_s P_r^s(z; x) \quad (55)$$

Therefore

$$\eta = \frac{M_\pi}{10T} \left(\frac{2T}{M_\pi} \right)^{5/2} A \int_0^\infty dx \frac{x^{5/2}}{z^{-1} e^x - 1} \sum_{s=0}^{\infty} B_s P_r^s(z; x) \quad (56)$$

and the distribution function as

$$f = f^0 + \frac{f^0}{T} \left(V_{ij} - \frac{1}{3} \delta_{ij} V_{hh} \right) \left(v_i v_j - \frac{1}{3} \delta_{ij} v^2 \right) \sum_{s=0}^{\infty} B_s P_r^s \left(z; \frac{M_\pi}{2T} v^2 \right) \quad (57)$$

For all the families of polynomials, i.e., for all the r values, we have exact conservation of the collision invariants. This fact makes possible to choose r so that we get the simplest integral. The appropriate value turns out to be $r = 5/2$ and then we get

$$\eta = \frac{A\pi^{3/2}}{2} \left(\frac{2T}{M_\pi} \right)^{5/2} g_{7/2}(z) B_0 \quad (58)$$

In this way the computation of the viscosity of the pion gas has been reduced to the computation of the B_0 coefficient.

VI. SIMPLIFYING THE UEHLING-UHLENBECK EQUATION

As discussed above the perturbed distribution function must be a solution of the Uehling-Uhlenbeck equation (15). In order to solve this equation it is quite convenient to write the l.h.s. in terms of macroscopic quantities, as the temperature, velocity, density and fugacity, and their derivatives. This can be done by using the equation of state and the hydrodynamic equations as follows: First of all at lowest order we have

$$\partial_t f + \vec{v} \vec{\nabla} f \simeq \partial_t f^0 + \vec{v} \vec{\nabla} f^0 = A \partial_t \left(\frac{1}{z^{-1} \exp\left(\frac{E}{T}\right) - 1} \right) + A \vec{v} \vec{\nabla} \left(\frac{1}{z^{-1} \exp\left(\frac{E}{T}\right) - 1} \right) \quad (59)$$

Thus the temporal derivative is

$$\partial_t f^0 = - \frac{f^{02} z^{-1} \exp\left(\frac{E}{T}\right)}{A} \left(\frac{1}{T} \partial_t E - \frac{E}{T^2} \partial_t T - \partial_t \log z \right) \quad (60)$$

where the energy E is given by

$$E = \frac{M_\pi}{2} (\vec{v} - \vec{V})^2 \quad (61)$$

so that we have

$$\partial_t f^0 = \frac{f^{02}}{Az \exp(-\frac{E}{T})} \left(\partial_t \log z + \frac{E}{T^2} \partial_t T + \frac{M_\pi}{T} v_i \partial_t V_i \right) \quad (62)$$

In a similar way we can obtain

$$v_j \nabla_j f^0 = \frac{f^{02} v_j}{Az \exp(-\frac{E}{T})} \left(\nabla_j \log z + \frac{E}{T^2} \nabla_j T + \frac{M_\pi}{T} v_i \nabla_j V_i \right) \quad (63)$$

Thus we have

$$\partial_t f^0 + v_j \nabla_j f^0 = \frac{f^{02}}{Az \exp(-\frac{E}{T})} \left(\partial_t \log z + \frac{E}{T^2} \partial_t T + \frac{M_\pi}{T} v_i \partial_t V_i + v_j \nabla_j \log z + \frac{E}{T^2} v_j \nabla_j T + \frac{M_\pi}{T} v_i v_j V_{ij} \right) \quad (64)$$

Here it is possible to use the state equation for the free pion gas and the ideal fluid equations (Euler, continuity and energy conservation) to find

$$\partial_t f^0 + v_j \nabla_j f^0 = \frac{f^{02}}{Az \exp(-\frac{E}{T})} \left[\frac{1}{T} \left(\frac{E}{T} - \frac{5}{2} \frac{g_{5/2}(z)}{g_{3/2}(z)} \right) v_i \nabla_i T + \left(\frac{M_\pi}{T} v_i v_j - \frac{2}{3} \frac{E}{T} \delta_{ij} \right) V_{ij} \right] \quad (65)$$

On the other hand, substituting expression (37) into (16), expanding to first order in the perturbation χ and taking benefit from the equilibrium distribution function relation

$$f_1^{0'} f_2^{0'} (1 + A^{-1} f_1^0) (1 + A^{-1} f_2^0) = f_1^0 f_2^0 (1 + f_1^{0'}) (1 + A^{-1} f_2^{0'}) \quad (66)$$

we obtain

$$\frac{A^2}{T} \frac{z^{-2} e^{E/T}}{(z^{-1} e^{E_1/T} - 1) (z^{-1} e^{E_2/T} - 1) (z^{-1} e^{E'_1/T} - 1) (z^{-1} e^{E'_2/T} - 1)} \Delta \left[\chi \left(1 - z \exp \left(-\frac{E}{T} \right) \right) \right] \quad (67)$$

where $E = E_1 + E_2 = E'_1 + E'_2$ and the Δ symbol is defined as

$$\Delta[f(x)] \equiv f(x'_1) + f(x'_2) - f(x_1) - f(x_2) \quad (68)$$

Thus the whole collision term can be written as

$$C[f_1] = \frac{1}{A^2 T} \int d\vec{v}_2 d\sigma \, u \, z^{-2} \exp(E/T) f_1 f_2 f'_1 f'_2 \Delta \left[\chi \left(1 - z \exp \left(-\frac{E}{T} \right) \right) \right] \quad (69)$$

where we have omitted the superindex 0 in the equilibrium distribution function. In this way the complete transport equation for the slightly unhomogeneous pion gas becomes

$$\begin{aligned} \frac{f_1^2}{z e^{-\frac{M_\pi v_1^2}{2T}}} \left[\left(\frac{M_\pi v_1^2}{2T} - \frac{5}{2} \frac{g_{5/2}(z)}{g_{3/2}(z)} \right) v_{1i} \nabla_i T + M_\pi \left(v_{1i} v_{1j} - \frac{1}{3} v_1^2 \delta_{ij} \right) V_{ij} \right] = \\ \frac{1}{A} \int d\vec{v}_2 d\sigma \, u \, z^{-2} e^{\frac{M_\pi}{2T} (4U^2 + u^2)} f_1 f_2 f'_1 f'_2 \Delta \left[\chi \left(1 - z e^{-\frac{M_\pi v^2}{2T}} \right) \right] \end{aligned} \quad (70)$$

where we have defined $\vec{U} = (\vec{v}_1 + \vec{v}_2)/2 = (\vec{v}'_1 + \vec{v}'_2)/2$, i.e. it is the collision center of mass velocity. Once again $\vec{u} = \vec{v}_1 - \vec{v}_2$ is the relative velocity of the incident pions. Thus, in particular, we have

$$E = \frac{M_\pi}{2} \left(2U^2 + \frac{1}{2} u^2 \right) \quad (71)$$

VII. LOW-ENERGY PION CROSS-SECTION

In order to solve the above transport equation it is needed to know the differential cross section for pion scattering. As we are interested in the low energy regime the most appropriate approach is the Chiral Perturba-

tion Theory which relies in the chiral symmetry pattern of the strong interactions. In particular it is quite useful the chiral Lagrangian approach suggested by Weinberg and extended up to the one loop level by Gasser and Leutwyler. This chiral Lagrangian approach consists in

$$T_{a_1, a_2, a'_1, a'_2}(s, t, u) = \frac{s - M_\pi^2}{F_\pi^2} \delta_{a_1 a_2} \delta_{a'_1 a'_2} + \frac{t - M_\pi^2}{F_\pi^2} \delta_{a_1 a'_1} \delta_{a_2 a'_2} + \frac{u - M_\pi^2}{F_\pi^2} \delta_{a_1 a'_2} \delta_{a_2 a'_1} \quad (72)$$

where the standard Mandelstam variables s , t and u are related by $s + t + u = 4M_\pi^2$, the a subindices refer to the pion isospin and F_π is the pion decay constant. In the total isospin basis there are three independent amplitudes corresponding to $I = 0, 1, 2$ which are given by

$$\begin{aligned} T^0 &= \frac{2s - M_\pi^2}{F_\pi^2} \\ T^1 &= \frac{t - u}{F_\pi^2} \\ T^2 &= \frac{2M_\pi^2 - s}{F_\pi^2} \end{aligned} \quad (73)$$

The averaged modulo squared amplitude is given by

$$|T|^2 = \frac{1}{\sum_I (2I+1)} \sum_I (2I+1) |T^I|^2 \quad (74)$$

which leads to

$$|T|^2 = \frac{1}{9F_\pi^4} \left(21M_\pi^4 + 9s^2 - 24M_\pi^2 s + 3(t-u)^2 \right) \quad (75)$$

The relativistic differential cross-section for elastic pion scattering is

$$d\sigma = \frac{|T|^2}{4uE_1E_2} dLIPS \quad (76)$$

a systematic expansion on the pion field derivatives and the pion mass which are considered to be of the same order. From the lowest order of this expansion, the pion elastic scattering amplitude is given by

where the differential Lorentz Invariant Phase Space is given by

$$dLIPS \equiv C (2\pi)^4 \delta^4(p_1 + p_2 - p'_1 - p'_2) \frac{dp'_1}{2(2\pi)^3 E'_1} \frac{dp'_2}{2(2\pi)^3 E'_2} \quad (77)$$

where the constant $C = 1/2$ takes into account the identity of the final pions.

As discussed above we are interested in the non relativistic regime where $\vec{p}_i = M_\pi \vec{v}_i$, $\vec{P} = 2M_\pi \vec{U}$ and $E_i = M_\pi + M_\pi v_i^2/2$. In this limit it is convenient to write the cross section in terms of the center of mass and relative velocities $\vec{U}$, $\vec{u}$ and $\vec{u}'$ defined as

$$\vec{u}' = \vec{v}'_1 - \vec{v}'_2 \quad u = u' \quad (78)$$

After some standard calculations the non relativistic reduction of the cross section can be found to be

$$d\sigma = \frac{23}{768} \frac{M_\pi^2}{\pi F_\pi^4} \left(1 - \frac{3}{2} U^2 + \frac{13}{184} u^2 + U^2 \cos \theta' \frac{2U + u \cos \theta'}{2U \cos \theta' + u} \right) d \cos \theta' \quad (79)$$

where θ' is the angle between by $\vec{U}$ and $\vec{u}'$.

VIII. LOWEST ORDER VISCOSITY FOR THE PION GAS

As we have seen in previous sections the perturbation of the distribution function appropriate for the compu-

tation of the pion gas viscosity can be written as

$$\chi = - \left(V_{ij} - \frac{1}{3} \nabla_h V_h \delta_{ij} \right) \left(v_i v_j - \frac{1}{3} v^2 \delta_{ij} \right) \sum_{s=0}^{\infty} B_s P_{5/2}^s(z; x) \quad (80)$$

where the B_s coefficients must be obtained by solving the linearized transport equation being B_0 the dominant one. In order to do so let us consider for a while a tensor l_{kl} which is an arbitrary function on z and $\hat{v}_1$. For further convenience we have defined the hat on any velocity as

$$\hat{v} = \sqrt{\frac{M_\pi}{2T}} v \quad (81)$$

By multiplying this tensor by the transport equation and integrating on v_1 we have

$$\begin{aligned} \int d\vec{v}_1 l_{kl}(z; \hat{v}_1) \frac{f_1^2}{ze^{-\hat{v}_1^2}} \left[\left(\hat{v}_1^2 - \frac{5}{2} \frac{g_{5/2}(z)}{g_{3/2}(z)} \right) v_{1i} \nabla_i T + M_\pi \left(v_{1i} v_{1j} - \frac{1}{3} v_1^2 \delta_{ij} \right) V_{ij} \right] = \\ -V_{ij} \sum_{s=0}^{\infty} B_s \frac{1}{A} \int d\vec{v}_1 d\vec{v}_2 d\sigma l_{kl}(z; \hat{v}_1) u z^{-2} e^{(2\hat{U}^2 + \frac{1}{2}\hat{u}^2)} f_1 f_2 f'_1 f'_2 \Delta \left[\left(v_i v_j - \frac{1}{3} v^2 \delta_{ij} \right) \left(1 - ze^{-\hat{v}_1^2} \right) P_{5/2}^s(z; \hat{v}^2) \right] \end{aligned} \quad (82)$$

where we have assumed the divergence of the velocity to vanish since the pion gas is taken to be incompressible. Now we choose the tensor to be

$$l_{kl}(z; \hat{v}) = \left(v_k v_l - \frac{1}{3} v^2 \delta_{kl} \right) \left(1 - ze^{-\hat{v}^2} \right) P_{5/2}^t(z; \hat{v}^2) \quad (83)$$

Then, one of the two integrals appearing in the r.h.s. of the equation (82) vanishes, namely

$$\int d\vec{v}_1 l_{kl}(z; \hat{v}_1) \frac{f_1^2}{ze^{-\hat{v}_1^2}} \left(\hat{v}_1^2 - \frac{5}{2} \frac{g_{5/2}(z)}{g_{3/2}(z)} \right) v_{1i} \nabla_i T = 0 \quad (84)$$

Therefore, the two sides of the equation (82) are proportional to V_{ij} which is arbitrary. Thus one possible solution to the equation is found by eliminating this tensor in both sides and then by contracting with the tensors $\delta_{ik} \delta_{jl}$ so that the r.h.s. of the equation (82) becomes

$$\begin{aligned} A^2 V_{ij} \int d\vec{v} \frac{M_\pi (v_k v_l - \frac{1}{3} v^2 \delta_{kl})^2}{z^{-1} e^{\hat{v}^2} - 1} P_{5/2}^t(z; \hat{v}^2) = \\ V_{ij} A^2 \frac{5\pi^{3/2} M_\pi}{2} \left(\frac{2T}{M_\pi} \right)^{7/2} g_{7/2}(z) \delta_{0t} \end{aligned} \quad (85)$$

In order to compute the B_s coefficients we have to proceed in a similar way with the r.h.s. of the equation (82). For that purpose we will take the low energy limit of the differential cross section which is just

$$d\sigma = \frac{23M_\pi^2}{768\pi F_\pi^4} d\cos\theta' \quad (86)$$

where θ' belongs to the interval $(0, \pi)$. Therefore, at very low energies the elastic pion scattering cross section can be considered as a constant, so that it is formally equivalent to the elastic scattering of rigid spheres of radius

$$R = \frac{M_\pi}{\pi F_\pi^2} \sqrt{\frac{23}{384}} \quad (87)$$

After the manipulations described below the r.h.s of the transport equation can be written the collision term as

$$V_{ij} \frac{23M_\pi^2 A^3}{768F_\pi^4} \left(\frac{2T}{M_\pi} \right)^{11/2} \sum_{s=0}^{\infty} B_s b_{st}(z) \quad (88)$$

where we have defined

$$\begin{aligned} b_{st}(z) = \int_0^\infty dU du \int_0^\pi d\theta' d\theta \int_0^{2\pi} d\phi \sin\theta' \sin\theta U^2 u^3 z^{-2} e^{2U^2 + \frac{1}{2}u^2} \prod_{a_1, a_2, a'_1, a'_2} \left(z^{-1} e^{v_a^2} - 1 \right)^{-1} \\ \Delta \left[\left(v_k v_l - \frac{1}{3} \delta_{kl} v^2 \right) \left(1 - ze^{-v^2} \right) P_{5/2}^s(z; v^2) \right] \Delta \left[\left(v_k v_l - \frac{1}{3} \delta_{kl} v^2 \right) \left(1 - ze^{-v^2} \right) P_{5/2}^t(z; v^2) \right] \end{aligned} \quad (89)$$

Then the transport equation (82) can be written as

$$\frac{23M_\pi A}{1920\pi^{3/2} F_\pi^4} \left(\frac{2T}{M_\pi} \right)^2 g_{7/2}^{-1}(z) \sum_{s=0}^{\infty} B_s b_{st} = \delta_{0t} \quad (90)$$

Therefore by defining

$$\mathcal{B}_s \equiv \frac{23M_\pi A}{1920\pi^{3/2} F_\pi^4} \left(\frac{2T}{M_\pi} \right)^2 g_{7/2}^{-1}(z) B_s \quad (91)$$

the viscosity is given by

$$\eta = \sqrt{\frac{2T}{M_\pi}} \frac{960\pi^3 F_\pi^4}{23M_\pi} g_{7/2}^2(z) \mathcal{B}_0 \quad (92)$$

where $\mathcal{B}_0$ is a solution of the equation system

$$\sum_{s=0}^{\infty} \mathcal{B}_s b_{ts}(z) = \delta_{t0} \quad (93)$$

or

$$\begin{pmatrix} b_{00}(z) & b_{01}(z) & b_{02}(z) & \cdots \\ b_{10}(z) & b_{11}(z) & b_{12}(z) & \cdots \\ b_{20}(z) & b_{21}(z) & b_{22}(z) & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} \mathcal{B}_0 \\ \mathcal{B}_1 \\ \mathcal{B}_2 \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ \vdots \end{pmatrix} \quad (94)$$

In order to find the leading contribution to the viscosity we can truncate the above system to find

$$\mathcal{B}_0 = \frac{1}{b_{00}(z)} \quad (95)$$

In this way the calculation of the viscosity have been reduced to the computation of the integral $b_{00}(z)$.

According to our previous discussion the viscosity of the pion gas at low density and temperatures is given by equation (92), where $\mathcal{B}_0$ is the solution of the equation system given in (93) and the corresponding coefficients are defined in (89). Thus it is possible, at least in principle, to find the viscosity as a function of the temperature T and the fugacity z (or the pion chemical potential). Alternatively it is also possible to find the viscosity as a function of the temperature and the pion number density. Due to the complicated integrals appearing in the definition of the $b_{st}(z)$ functions it has been possible to get numerical results only by Monte Carlo [12] integration. For the sake of simplicity most of these numerical results have been obtained by using the approximation of (95) and the hard sphere cross section of (86). In order to check the validity of these approximations we have computed the 4×4 matrix containing the first $b_{st}(z)$ functions for the particular value $z = 1$. The result and the corresponding errors is

$$b_{st} = \begin{pmatrix} \begin{array}{c|c|c|c} 277.2 \pm 0.7 & -42 \pm 1 & -99 \pm 3 & -408 \pm 14 \\ \hline -42 \pm 1 & 1043 \pm 8 & -395 \pm 8 & -800 \pm 30 \\ \hline -99 \pm 3 & -395 \pm 8 & 10700 \pm 110 & -7100 \pm 200 \\ \hline -408 \pm 14 & -800 \pm 30 & -7100 \pm 200 & 199000 \pm 2000 \end{array} \end{pmatrix} \quad (96)$$

Now we consider the submatrices 1×1 , 2×2 , 3×3 and 4×4 and solve the corresponding system for the appropriate number of $\mathcal{B}$ so that we can check the convergence of the results as the size of the truncation is increased. The results are shown in table 1. There we see that the error is of the order of 1.6 %

	$\mathcal{B}_0$	$\mathcal{B}_1$	$\mathcal{B}_2$	$\mathcal{B}_3$
Zero order	$3.608 \cdot 10^{-3}$			
First order	$3.630 \cdot 10^{-3}$	$1.497 \cdot 10^{-4}$		
Second order	$3.647 \cdot 10^{-3}$	$1.654 \cdot 10^{-4}$	$3.985 \cdot 10^{-5}$	
Third order	$3.666 \cdot 10^{-3}$	$1.765 \cdot 10^{-4}$	$4.700 \cdot 10^{-5}$	$9.902 \cdot 10^{-6}$

To take into account the velocity effects in the cross section we will consider (79) instead of (86) in the computation of the b_{ij} integrals. Then we will have

$$b_{st} = b_{st}^0 + \frac{T}{M_\pi} b_{st}^1 \quad (97)$$

where b_{ij}^0 is the leading term corresponding to the hard sphere approximation and

$$b_{00}^1 = \int_0^\infty dU du \int_0^\pi d\theta d\alpha \int_0^{2\pi} d\phi \frac{\sin \theta \sin \alpha U^2 u^3 e^{2U^2 + \frac{1}{2}u^2}}{\left(e^{v_1'^2} - 1\right) \left(e^{v_2'^2} - 1\right) \left(e^{v_1^2} - 1\right) \left(e^{v_2^2} - 1\right)} \left(\frac{13}{92} u^2 - 3U^2 + 2U^2 \cos \theta \frac{2U + u \cos \theta}{2U \cos \theta + u} \right) \left(\Delta \left[\left(v_i v_j - \frac{1}{3} \delta_{ij} v^2 \right) \left(1 - e^{-v^2} \right) \right] \right)^2 \quad (98)$$

At the zero order in the sense of the previous table we have

$$\mathcal{B}_0 = \frac{1}{b_{00}} = \frac{1}{b_{00}^0 + \frac{T}{M_\pi} b_{00}^1} \quad (99)$$

which at low temperatures can be written as

$$\mathcal{B}_0 = \frac{1}{b_{00}^0} \left(1 - \frac{T}{M_\pi} \frac{b_{00}^1}{b_{00}^0} \right) \quad (100)$$

so that

$$\eta = \sqrt{\frac{2T}{M_\pi}} \frac{960\pi^3 F_\pi^4}{23M_\pi} g_{7/2}^2(z) \frac{1}{b_{00}^0} \left(1 - \frac{T}{M_\pi} \frac{b_{00}^1}{b_{00}^0} \right) \quad (101)$$

This formula can be applied provided

$$T \ll M_\pi \left| \frac{b_{00}^0}{b_{00}^1} \right| \quad (102)$$

After a numerical integration we find, for $z = 1$, $b_{00}^1 = -310 \pm 40$. By comparison with our previous result $b_{00}^0 = 277.2 \pm 0.7$ we get the upper bound for the temperature

$$T \ll (0.90 \pm 0.12) M_\pi \quad (103)$$

which means that the hard sphere approximation is safe provided the temperature is much smaller than the 90% of the pion mass. As we are working in a non relativistic framework this will always be the case and then no additional constrain on the applicability of our results is coming from the hard sphere approximation.

Therefore in the following we will work at zero order and in the hard sphere approximation. In this case the viscosity is given by

$$\eta(T, z) = \sqrt{T} \frac{4096\sqrt{2}\pi^2 F_\pi^4}{345M_\pi^{3/2}} (b_{00}(z))^{-1} \left[\int_0^\infty \frac{x^{5/2}}{z^{-1}e^x - 1} dx \right]^2 \quad (104)$$

Some numerical results obtained from this formula are found in table 2 for different values of the fugacity z .

	$\eta \text{ (MeV}^3\text{)}$
Quantum case ($z = 1$)	$\sqrt{T} \ 4.00 \cdot 10^5$
Quantum case ($z = 0.05$)	$\sqrt{T} \ 1.40 \cdot 10^6$
Classical case	$\sqrt{T} \ 1.9 \cdot 10^6$

The classical case is just the limit of z going to zero. This corresponds to a classical gas of hard spheres of radius

$$R = \frac{M_\pi}{\pi F_\pi^2} \sqrt{\frac{23}{384}} \quad (105)$$




FIG. 1: Plot of $(b_{00}(z))^{-1} \left[\int_0^\infty \frac{x^{5/2}}{z^{-1}e^x - 1} dx \right]^2$ versus z . It can be understood as a plot of the viscosity, for constant temperature and in arbitrary units, versus the fugacity.

Because of the particular way we have performed our computations the particular case $z = 0$ is not numerically accessible. However this case has been treated long time ago and it is possible to find an analytic result for the viscosity, namely

$$\eta = \sqrt{T} \frac{5\sqrt{M_\pi}}{16\sqrt{\pi}R^2} \quad (106)$$

Thus we can check that our computations have the proper $z = 0$ limit. In fig.1 we show the behaviour of b_{00} and the viscosity, for constant T , in terms of the fugacity. From this plots we learn that the viscosity of the bosonic quantum gas is smaller than that of the classical gas. This could be expected because of the following heuristic argument: After some elastic collision the emerging particles have more affinity for occupied states than classical ones. Therefore the microscopic transfer of momentum is more effective in a classical gas than in a bosonic gas and then the viscosity is also larger. A pictorial view of this fact is displayed in fig.2.

In many cases it is useful to have the viscosity as a function of the temperature and the density instead of the temperature and the fugacity. The pionic density can be written in terms of the temperature and the fugacity as

$$\frac{n}{T^{3/2}} = \frac{gM_\pi^{3/2}}{\sqrt{2}\pi^2} \int_0^\infty \frac{x^{1/2}}{z^{-1}e^x - 1} dx \quad (107)$$

This equation defines also implicitly the fugacity as a function of the density and the temperature $z = z(n, T)$. However, due to the complexities of the above integral this functions have only been computed numerically. In fig.4 we plot $\eta/\sqrt{T}$ as a function of $n/T^{3/2}$. The computed points can be fitted quite well in the plotted range with a polynomial

$$f_1(x) = A + Bx + Cx^2 \quad (108)$$




FIG. 2: *a)* Two particles, belonging to two different regions of the gas with different average momentum (black and white), before an elastic collision. *b)* Particles before the collision in the classical case where they typically get a new momentum (represented in grey) *c)* Particles in the bosonic gas where they typically want to go to occupied states with the same momentum.

with constants $A = 0.1814 \pm 0.0004$, $B = (-2.42 \pm 0.03) \cdot 10^{-4}$ and $C = (9.6 \pm 0.3) \cdot 10^{-8}$. Thus the function obtained (fig.3) for the viscosity versus the temperature and the density is

$$\eta(T, n) = \frac{4096\sqrt{2}\pi^2 F_\pi^4}{345M_\pi^{3/2}} \sqrt{T} \left[0.1814 - 2.42 \cdot 10^{-4} \frac{n}{T^{3/2}} + 9.6 \cdot 10^{-8} \left(\frac{n}{T^{3/2}} \right)^2 \right] \quad (109)$$




FIG. 3: Numerical fit of $\eta/\sqrt{T} \propto f_1(n/T^{3/2})$.

As it may be noticed in fig.4, the values of the viscosity cover the range from 10^6 to 10^7 MeV³ for the considered densities and temperatures.

X. CONCLUSIONS

In this work we have computed the viscosity of the pion gas, starting from first principles only. This computation

is relevant for the hydrodynamic description of hadronic matter at low energies and densities. The work relies on the use of the Uehling-Uhlenbeck equation, which is the quantum version of the Boltzmann equation, and chiral symmetry, which determines completely the structure of the pion interactions at low energies. We have also found a formula for the viscosity in terms of the temperature and the density which properly fits our numerical results. The interest of the computations is twofold. First it is useful to check when the usual assumption of perfect fluid for the hadronic fluid is reasonable enough. Second it can be used to include the viscosity in the Navier-Stokes equations when dissipative effects cannot be neglected. The main limitation of our work is that it can be applied only in the non-relativistic domain. As a consequence of that it cannot be used directly in the study of the events observed in the modern ultra relativistic heavy ion colliders such as RHIC or the future LHC. In any case we understand that our work is interesting in order to show that the computations of transport coefficients of the pion gas can be done in a complete model independent way. In fact we consider that the result presented here is just the first step of a complete relativistic computation which could be applied at higher energies and therefore in more realistic situations, to improve the standard hydrodynamic description of hadronic matter. Work is in progress in that direction.




FIG. 4: 3D plot of viscosity vs. temperature and particle number density, given by equation (109).

Acknowledgments

The authors want to thank A. Gomez Nicola and J.R. Pelaez for reading the manuscript. This work has been partially supported by the Ministerio de Educación y Ciencia (Spain) (CICYT AEN 97-1693 and PB98-0782).

XI. APPENDIX

In this appendix we study the main properties of the family of orthogonal polynomials $P_r^s(z; x)$ defined on the

interval $(0, \infty)$ with measure

$$d\mu_r(z; x) \equiv \frac{x^r}{z^{-1}e^x - 1} \quad (110)$$

where $z \in (0, 1]$ and $r > 0$. By definition the polynomials are orthogonal for some given r , or in other words,

$$\begin{aligned} (P_r^s, P_r^{s'}) &= \int_0^\infty dx \frac{x^r}{z^{-1}e^x - 1} P_r^s(z; x) P_r^{s'}(z; x) \\ &= A(z; r, s) \delta_{ss'} \end{aligned} \quad (111)$$

For simplicity we define the polynomials so that they are monic, i.e. the coefficient of the term of highest degree in each polynomial is taken to be one. Thus, the first polynomial is always the unity and his norm is given by

$$\begin{aligned} P_r^0 = 1 \rightarrow (P_r^0, P_r^0) &= \int_0^\infty dx \frac{x^r}{z^{-1}e^x - 1} \\ &= \Gamma(r+1) g_{r+1}(z) \end{aligned} \quad (112)$$

From the condition

$$(P_r^0, P_r^1) = 0 \quad (113)$$

it is possible to compute the second polynomial which turns out to be

$$P_r^1(z; x) = \frac{g_{r+2}(z)}{g_{r+1}(z)} (r+1) - x \quad (114)$$

The third polynomial is obtained from the conditions $(P_r^0, P_r^2) = 0$ and $(P_r^1, P_r^2) = 0$ which gives

$$\begin{aligned} P_r^2(z; x) &= \frac{(r+3)g_{r+2}(z)g_{r+4}(z) - (r+2)g_{r+3}^2(z)}{(r+2)g_{r+1}(z)g_{r+3}(z) - (r+1)g_{r+2}^2(z)} (r+2)(r+1) + \\ &\quad \frac{(r+3)g_{r+1}(z)g_{r+4}(z) - (r+1)g_{r+2}(z)g_{r+3}(z)}{(r+1)g_{r+2}^2(z) - (r+2)g_{r+1}(z)g_{r+3}(z)} (r+2)x + x^2 \end{aligned} \quad (115)$$

Higher polynomials can be obtained in a similar way.

-
- [1] J. D. Bjorken. *Phy. Rev.* **D27** 140(1983)
 - [2] L. D. Landau, *Izv. Akad. Nauk. SSSR D* **17**, 51 (1953)
 - [3] P. Gerber, H. Leutwyler and J.L. Goity *Phy. Let.* **B246** 513 (1990)
 - [4] S. Weinberg, *Physica* **96A** 327 (1979)
J. Gasser and H. Leutwyler, *Ann. of Phys.* **158** 142

-
- (1984)
 - [5] E.M. Lifshitz, L.P. Pitaevskii, *Kinetic Theory*, Volume 10 *Course of theoretical physics*, Pergamon Press (1981)
 - [6] A. Dobado and J.R. Peláez. *Phy. Rev.* **D59** 034004(1999)
 - [7] D.J. Amit, Y. Verbin, *Statistical physics, an introductory course*, World scientific (2000).

- [8] M. Abramowitz, I.A. Stegun, *Handbook of mathematical functions*, Dover (1968).
- [9] R.L. Liboff, *Kinetic theory, classical, quantum and relativistic descriptions*, Prentice Hall (1999)
- [10] E.A. Uehling, G.E. Uhlenbeck, *Phys. Rev.* **43** 552 (1933).
- [11] R. Balescu, *Equilibrium and nonequilibrium statistical mechanics*, John Wiley (1975)
- [12] W.H. Press et al. *Numerical recipes in C*, Cambridge Univ. Press (1997).